Blueberries



Sinbar[®]WDG

Agricultural Herbicide



New! WDG Formulation is Easier and Faster to Mix and Load

The new Sinbar WDG formulation is free-flowing, virtually dust-free and easier to handle, measure and mix.



TESSENDERLO KERLEY, INC. 2255 North 44th Street Suite 300 Phoenix, AZ 85008-3279

Customer Service: 1-800-525-2803 1-602-889-8300

www.novasource.com

Application of Sinbar WDG Agricultural Herbicide for Blueberries in the U.S.A. and Canada

Sinbar[®] WDG is a dependable tool for use as a pre-emergent herbicide in blueberry production. Use Sinbar WDG in a rotation or tank mix program. Do not use Sinbar WDG on blueberries established for less than one year. Sinbar WDG controls susceptible weeds for an extended period of time. The degree of control will be based on many factors including:

Resistance Management

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally occurring resistant biotypes may survive, propagate and become dominant in that field.

To manage herbicide resistance, change cultural practices within and between crop seasons such as using a combination of tillage, re-treatment, tank-mix partners and/ or sequential herbicide applications that have a different mode of action.

Proper Application Rates and Techniques

Consult the Sinbar WDG specimen label for complete instructions, precautions and recommendations. For a copy of the specimen label go to *www.novasource.com*. Consult local agricultural experts for specific recommendations available in your area.

Soil Texture and Soil Organic Matter

Sinbar WDG is normally not for use on sand and gravel soil types. Where a range of dosages is given, use the lower rate on coarse-textured soils (low in clay or organic matter) and the higher rate on fine-textured soils (high in clay or organic matter). Care must be taken for use of Sinbar WDG on soils with less than one percent organic matter, as severe crop injury may occur. Refer to the label for full use directions.

Soil Moisture and Soil Temperature

Moisture is required to activate the chemical; best results occur if rainfall (or sprinkler irrigation) occurs within 2 weeks after application.

Warm temperatures accelerate the translocation of Sinbar WDG. Weed control may be slower at lower temperatures.

Important: Always read the label before buying and follow label instructions when using this product.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Tessenderlo Kerley, Inc. warrants only that this product conforms to the product description on the label. Except as warranted by this label, Tessenderlo Kerley, Inc. makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. Tessenderlo Kerley, Inc. does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, Tessenderlo Kerley, Inc.'s maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability imitations, buyer may return the unopened container to the place of purchase for full refund. Buyer's use of this product shall constitute conclusive evidence of buyer's acknowledgement and acceptance of the forgoing limitations. Some states do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller's sales invoice for this product.

Sinbar is a registered trademark of Tessenderlo Kerley, Inc. ©2006, 2008 Tessenderlo Kerley Inc. All rights reserved. NS PIUS Sinbar WDG Blueberry v16 10/11